arXiv:1212.1899v1 [math.CO] 9 Dec 2012

Extensions of Sperner and Tucker's lemma for manifolds

Oleg R. Musin[*]

Abstract

The Sperner and Tucker lemmas are combinatorial analogous of the Brouwer and Borsuk - Ulam theorems with many useful applications. These classic lemmas are concerning labellings of triangulated discs and spheres. In this paper we show that discs and spheres can be substituted by large classes of manifolds with or without boundary.

Keywords: Sperner's lemma, Tucker's lemma, the Borsuk-Ulam theorem.

1 Introduction

Sperner's lemma is a statement about labellings of triangulated simplices (d-balls). It is a discrete analog of the Brouwer fixed point theorem.

Let T be a triangulation of a PL manifold M^d. Suppose that each vertex of T is assigned a unique label from the set $\{1, 2, \dots, n\}$. Such a labelling is called a *n-labelling* of T. We say that a d-simplex in T is a *fully labelled simplex* or simply a *full cell* if all its labels are distinct.

Sperner's lemma [13] states:
Every Sperner labelling of a triangulation of an d-dimensional simplex contains a cell labelled with a complete set of labels: $\{1, 2, \dots, d+1\}$.

There are several extensions of this lemma. One of the most interesting is the De Loera - Petersen - Su theorem. In [3] they proved the Atanassov conjecture [1]:
Let P be a d-dimensional polytope with vertices $v_1, \dots, v_n$. Let T be a triangulation of P. Let $L : V(T) \to \{1, 2, \dots, n\}$ be a (Sperner) labelling such that $L(v_i) = i$ and if $v \in V(T)$ lies in a face F of P, then $L(v) = i$, where v_i is one of the vertices of F.

Then there are at least $n - d$ fully-colored (i.e. with distinct labels) d-simplices of T.

Meunier [7] extended this theorem:
Let P^d be an embedded in $\mathbb{R}^d$ d-dimensional PL manifold with boundary B. Suppose B has n vertices $v_1, \dots, v_n$. Let T be a triangulation of P. Any Sperner labelling of T contains at

---

[*] This research is supported by the Russian government project 11.G34.31.0053 and NSF grant DMS - 1101688.

least $n + \lceil \min_i(\deg_B(v_i))/d \rceil - d - 1$ *fully-labelled* d*-simplices such that any pair of these fully-labelled simplices receives two different labellings.*

Here $deg_B(v)$ is the number of edges of B which v belongs to.

A discrete version of the Borsuk-Ulam theorem is *Tucker's lemma* [14]:
Let T *be an antipodal triangulation of the* d*-sphere* $\mathbb{S}^d$*. Let*

$$L : V(T) \to \{+1, -1, +2, -2, \dots, +d, -d\}$$

be an equivariant (or Tucker's) labelling, i.e. $L(-v) = -L(v)$*). Then there exists a complementary edge in* T *such that its two vertices are labelled by opposite numbers* (see [6, Theorem 2.3.1]).

This lemma was extended by Ky Fan [4]:
Let T *be an antipodal triangulation of* $\mathbb{S}^d$*. Suppose that each vertex* v *of* T *is assigned a label* $L(v)$ *from* $\{\pm1, \pm2, \dots, \pm n\}$ *in such a way that* $L(-v) = -L(v)$*. Suppose this labelling does not have complementary edges. Then there are an odd number of* d*-simplices of* T *whose labels are of the form* $\{k_0, -k_1, k_2, \dots, (-1)^d k_d\}$*, where* $1 \leq k_0 < k_1 < \dots < k_d \leq n$*. In particular,* $n \geq d + 1$*.*

In this paper we consider extensions of the Sperner, De Loera - Petersen - Su, Tucker and Fan theorems for manifolds. We show that for all cases d-balls and spheres can be substituted by d-manifolds with or without boundary.

2 Extensions of Sperner's lemma for manifolds

Throughout this paper we consider piecewise linear (PL) d-dimensional manifolds M^d. M is called a *PL manifold* if it is a topological manifold together with a piecewise linear structure on it. Every PL manifold M admits a triangulation: that is, we can find a collection of simplices T of dimensions $0, 1, \dots, d$, such that (1) any face of a simplex belonging to T also belongs to T, (2) any nonempty intersection of any two simplices of T is a face of each, and (3) the union of the simplices of T is M.

We consider here only two types of manifolds: closed or compact manifolds with boundaries. (A manifold M is called *closed* if it is a compact manifold without boundary.)

Definition. Let M^d be a closed d-dimensional PL manifold with vertices $V = \{v_1, \dots, v_n\}$ and faces $\{F_i\}$ of dimension from 1 to d. Let T be a triangulation of M such that for any face F_i it is a triangulation of F_i. Suppose that the vertices of T have a labelling satisfying the following conditions: each vertex v_k of V is assigned a unique label from $\{1, 2, \dots, n\}$, and each other vertex v of T belonging to a face F_i with vertices $V(F_i) := \{v_{i_1}, \dots, v_{i_m}\}$ from V is assigned a label of one of the vertices of $V(F_i)$. Such a labelling is called a *Sperner labelling* of T.

Let P be a set of n points $p_1, \dots, p_n$ in $\mathbb{R}^d$. Denote by $S(P)$ the collection of all simplices spanned by vertices $\{p_{i_1}, \dots, p_{i_k}\}$ with $1 \leq k \leq d + 1$. Consider a point $x \in \mathbb{R}^d$ and the set

$S_x(P)$ of all simplices from $S(P)$ which cover x. In the case if no such simplices we write $S_x(P) = \emptyset$. Let us remove from $S_x(P) \neq \emptyset$ all simplices such that its boundary contain x. Denote this set of simplices by $\mathrm{cov}_P(x)$ or just by $\mathrm{cov}(x)$.

Definition. Let $P := \{p_1, \dots, p_n\}$ be points in $\mathbb{R}^d$. Let T be a triangulation of a closed PL manifold M^m. Let L be an *n-labelling* of T, i.e. a labelling (map) $L : V(T) \to \{1, 2, \dots, n\}$. If for $v \in V(T)$ we have $L(v) = i$, then set $f_{L,P}(v) := p_i$. Therefore, $f_{L,P}$ is defined for all vertices of T. Then it uniquely defines a simplicial (piecewise linear) map $f_{L,P} : M^m \to \mathbb{R}^d$.

Theorem 2.1. *Let $P := \{p_1, \dots, p_n\} \subset \mathbb{R}^d$. Suppose $x \in \mathbb{R}^d$ is such that $\mathrm{cov}_P(x)$ consists of d-simplices. Then any n-labelling L of a triangulation T of a closed PL manifold M^d must contain an even number of full cells which are labelled as simplices in $\mathrm{cov}_P(x)$.*

Proof. Actually, a proof easily follows from the well known fact that the degree of a continues map of a closed manifold to a manifold is a topological invariant modulo 2 (see, for instance, [9]). Indeed, if $y \in \mathbb{R}^d$ lies outside of the convex hull of P in $\mathbb{R}^d$, then $f^{-1}_{L,P}(y) = \emptyset$. Therefore, for any point x in $\mathbb{R}^d$ which is a regular value of $f_{L,P}$, we have $|f^{-1}_{L,P}(x)| \equiv |f^{-1}_{L,P}(y)| \equiv 0 \pmod 2$. Thus, the number of full cells which are labelled as simplices in $\mathrm{cov}_P(x)$ is even. □

For the classical case $n = d + 1$ we have the following result (see also [5]):

Corollary 2.1. *Let T be a triangulation of a closed PL manifold M^d. Any $(d + 1)$-labelling of T must contain an even number of full cells.*

Corollary 2.2. *Let M^d be a compact PL manifold with boundary B. Let B be PL homeomorphic to the boundary of a d-simplex (i.e. $B \cong \mathbb{S}^{d-1}$). Then any $(d + 1)$-labelling of a triangulation T of M which is a Sperner labelling on the boundary B must contain an odd number of full cells; in particular, there is at least one.*

Proof. We prove this corollary by induction. It is clear for $d = 1$. Let S denotes a d-simplex. Two manifolds M and S can be glued together along B. We denote the new manifold by N. Corollary 2.1 implies that any $(d + 1)$-labelling of any triangulation of N has an even number of full cells.

Let us add to vertices of S one more vertex v which is an internal point of S. Let C be the "cone triangulation" of S with vertex v. Set $L(v) := d + 1$. So L is defined on the triangulation $T \cup C$ of N and has an even number of full cells.

Let F denotes the face of B with vertices $\{1, 2, \dots, d\}$. Then σ is a full cell of C only if its facet on B is a full cell in F. By induction, T on F contains an odd number of full cells. Thus, T has an odd number of full cells. □

Note that for the case when M is a d-ball Corollary 2.2 is Sperner's lemma.

It is clear, for the case when P is a d-simplex this theorem is Theorem 3.1. Now we show how from Theorem 2.1 follows the De Loera - Peterson - Su theorem.



Figure 1: Sperner's lemma for the Möbius strip

Corollary 2.3. *Let X be the vertex set of a convex polytope P in $\mathbb{R}^d$ with n vertices. Let M^d be a compact PL manifold with boundary P. Suppose $p \in \mathbb{R}^d$ is such that $\operatorname{cov}_X(p) \neq \emptyset$. Then any n-labelling of a triangulation of M which is a Sperner labelling on the boundary must contain an odd number of full cells which are labelled as simplices in $\operatorname{cov}_X(p)$.*

Proof. This corollary can be proved by the same arguments as Corollary 2.2. □

Corollary 2.4. *Let P be a convex polytope P in $\mathbb{R}^d$ with n vertices. Let M^d be a compact PL manifold with boundary B which is PL homeomorphic to P. Then any n-labelling of a triangulation of M which is a Sperner labelling on the boundary contains at least $n-d$ full cells.*

If M is a d-ball, then this statement is the De Loera - Peterson - Su theorem [3, Th. 1].

Proof. Actually, a proof of this corollary follows from another theorem from [3, Th. 4]: *Any convex polytope P in $\mathbb{R}^d$ with n vertices contains a pebble set of size $n-d$.* (A finite set of points (*pebbles*) in P is called a *pebble set* if each d-simplex of P contains at most one pebble interior to chambers.)

Consider a pebble set $\{p_i\}$ of size $n-d$. Then for $i \neq j$ we have $\operatorname{cov}(p_i) \cap \operatorname{cov}(p_j) = \emptyset$. Thus Corollary 2.3 guarantee that there are at least $n-d$ full cells. □

Remark. In fact, Meunier's proof of his extension of the De Loera - Peterson - Su theorem is not based on the "pebbles set theorem". It is an interesting problem to find an extension of this theorem and to derive Meunier's theorem for manifolds from Theorem 2.1.

3 Extensions of Tucker's lemma for manifolds

Let M be a closed PL-manifold with a free simplicial involution $T : M \to M$, i.e. $T^2(x) = x$ and $T(x) \neq x$ for all $x \in M$. For any $\mathbb{Z}_2$-manifold (M, T) we say that a map $f : M^m \to \mathbb{R}^d$ is *antipodal* (or equivariant) if $f(T(x)) = -f(x)$.

Definition. We say that a closed PL free $\mathbb{Z}_2$-manifold (M^d, T) is a *BUT (Borsuk-Ulam Type) manifold* if for any continuous $g : M^d \to \mathbb{R}^d$ there is a point $x \in M$ such that $g(T(x)) = g(x)$. Equivalently, if a continuous map $f : M^d \to \mathbb{R}^d$ is antipodal, then the zeros set $Z_f := f^{-1}(0)$ is not empty.

In [11] we found several equivalent necessary and sufficient conditions for manifolds to be BUT. For instance, *M^d is a BUT manifold if and only if M admits an antipodal continuous transversal to zeros map $h : M^d \to \mathbb{R}^d$ with $|Z_f| = 2 \pmod 4$.*

Let M^d be a closed PL manifold with a free involution T. Let Λ be any equivariant triangulation of M. We say that

$$L : V(\Lambda) \to \{+1, -1, +2, -2, \dots, +d, -d\}$$

is an *equivariant* (or *Tucker's*) labelling if $L(T(v)) = -L(v))$.

An edge e in Λ is called *complementary* if its two ends are labelled by opposite numbers, i.e. if $e = uv$, then $L(v) = -L(u)$.

Theorem 3.1. *A closed PL free $\mathbb{Z}_2$-manifold M^d is BUT if and only if for any equivariant labelling of any equivariant triangulation of M there exists a complementary edge.*

For the case $M = \mathbb{S}^d$ this is Tucker's lemma.

Proof. Any equivariant labelling L of a triangulation Λ of M defines a simplicial map $f_L : \Lambda \to C^d$, where C^d is the crosspolytope in $\mathbb{R}^d$ with vertex set $\{+1, -1, +2, -2, \dots, +d, -d\}$ [6, Sec. 2.3]. Indeed, let $e_1, \dots, e_d$ be an orthonormal basis of $\mathbb{R}^d$. We define $f_L : \Lambda \to \mathbb{R}^d$ for $v \in V(\Lambda)$ by $f_L(v) = e_i$ if $L(v) = i$ and $f_L(v) = -e_i$ if $L(v) = -i$. In other words, $f_L = f_{L,C}$ (see Section 2).

It is easy to see that if L has no complementary edges, then $f_L : \Lambda \to \mathbb{R}^d$ has no zeros.

Actually the reverse implication can be proved by the same arguments as equivalence of the Borsuk-Ulam theorem and Tucker's lemma in [6, 2.3.2], i.e. if there is continuous antipodal $f : M^d \to \mathbb{S}^{d-1}$ (i.e. $Z_f = \emptyset$) then can be constructed Λ and L with no complementary edges. (See also Theorem 3.2.) □

Theorem 3.1 and [11, Theorem 2] immediately imply:

Corollary 3.1. *Let M^d be a closed PL manifold with a free involution T. Then M is a BUT manifold if and only if there exist an equivariant triangulation Λ of M and an equivariant labelling of $V(\Lambda)$ such that $f_L : \Lambda \to \mathbb{R}^d$ is transversal to zeros and the number of complementary edges is $4k + 2$, where $k = 0, 1, 2, \dots$.*



Figure 2: Tucker's lemma for the Möbius strip

Corollary 3.2. *Let T be a triangulation of a PL compact manifolds M^d with the boundary $\mathbb{S}^{d-1}$. Assume Λ is antipodally symmetric on the boundary. (That means that the subset of simplices of Λ which are in $\mathbb{S}^{d-1}$ provides a triangulation od $\mathbb{S}^{d-1}$ where if s is a simplex then so $-s$.) Let $L : V(\Lambda) \to \{+1, -1, +2, -2, \dots, +d, -d\}$ be a labelling of the vertices of T which satisfies $L(-v) = -L(v)$ for all vertices v in $\mathbb{S}^{d-1}$. Then there is a complementary edge in Λ.*

Proof. Consider two copies of M: M_+ and M_-, where for M_+ we take a given labelling L and for M_- we take a labelling $\bar{L} = (-L)$, i.e. $\bar{L}(v) = -L(v)$. Since L is antipodal on the boundary $\mathbb{S}^{d-1}$ the connected sum $N := M\#M$ with a free involution $I : N \to N$, where $I(M_+) = M_-$, is well defined. Since [11, Corollary 1] implies that N is BUT, from Theorem 3.1 we have that there is a complementary edge. □

Now we extend Theorem 3.1 for n-labelling.

Theorem 3.2. *Let P be a centrally symmetric set of $2n$ points in $\mathbb{R}^d$. Let points in P be equivariantly labelled by $\{+1, -1, +2, -2, \dots, +n, -n\}$. Let M^d be a closed PL manifold with a free involution. Then M is a BUT manifold if and only for any equivariant triangulation Λ of M and for any equivariant labelling $L : V(\Lambda) \to \{+1, -1, +2, -2, \dots, +n, -n\}$ there exists a k-simplex s in Λ, $0 \le k \le d$, which is labelled as a k-simplex in $\mathbb{R}^d$ with vertices in P and which is contained 0, i.e. $f^{-1}_{L,P}(0) \in s$.*

Proof. If M is BUT, then $f_{L,P}$ has zeros, i.e. there is a simplex s as required.

Suppose M is not BUT. Then there is a continuous antipodal $h : M^d \to \mathbb{S}^{d-1}$. Let Q denotes the boundary of the convex hull of P in $\mathbb{R}^d$. Since Q is PL homemorphic to $\mathbb{S}^{d-1}$, h can be approximated by a PL antipodal $\tilde{h} : M^d \to Q$. This map gives a labelling L with $f_{L,P}^{-1}(0) = \emptyset$, a contradiction. $\square$

4 Radon partitions and Fan's lemma for manifolds

In this section we show that Ky Fan's lemma for manifolds follows from Theorem 3.2.

Radon's theorem on convex sets states that any set S of $d+2$ points in $\mathbb{R}^d$ can be partitioned into two (disjoint) sets A and B whose convex hulls intersect. Moreover, if $\operatorname{rank}(S) = d$, then this partition is unique. The partition $S = A \bigcup B$ is called the *Radon partition* of S.

Breen [2] proved that if S is a $(d{+}2)$-subset of the moment curve C_d in $\mathbb{R}^d$, then $S = A \bigcup B$ is the Radon partition if and only if A and B alternate along C_d. Actually, Breen's theorem can be extended for convex curves in $\mathbb{R}^d$.

We say that a curve K in $\mathbb{R}^d$ is *convex* if it intersects any hyperplane at no more than d points.

It is well known that the moment curve C_d is convex. In [10, Sec. 3] considered several other examples of convex curves.

Definition Let $K = \{x(t) = (x_1(t), \ldots, x_d(t)) : t \in [a, b]\}$. Let $S = \{x(t_1), \ldots, x(t_{d+2})\} \subset \mathbb{R}^d$, where $a < t_1 < t_2 < \ldots < t_{d+2} < b$. We say that A and B *alternate along* K if $S = A \bigcup B$ is the Radon partion and $A = \{x(t_1), x(t_3), \ldots\}$, $B = \{x(t_2), x(t_4), \ldots\}$.

Theorem 4.1. *A curve K in $\mathbb{R}^d$ is convex if and only if for any $(d+2)$-subset S of K its Radon partition sets A and B alternate along K.*

Proof. Let K be convex and $S = \{x(t_1), \ldots, x(t_{d+2})\}$ be a $(d+2)$-subset of K. Let $A \bigcap B$ be the Radon partition of S. In the case if A and B not alternate along K there is at most d points $P = \{x(\tau_i)\}$ which separate A and B on K. If $r = |P| < d$, we add to P $d - r$ points $x(\tau)$ with $\tau \in (a, t_1)$. Then P defines a hyperplane H passes through the points in P. Clearly, H separates A and B in $\mathbb{R}^d$. Thus, $A \bigcap B$ cannot be the Radon partition of S, a contradiction.

Suppose for any $(d+2)$-subset S of K its Radon partition sets A and B alternate along K. If K is not convex, then there is a hyperplane H which intersects K at $r \geq d+1$ points. Therefore, H separates K into $r+1$ connected components $C_1, \ldots, C_{r+1}$. Let $S = \{x(t_1), \ldots, x(t_{d+2})\}$, where $x(t_i) \in C_i$. Since A and B which alternate along K are separated by H the partition $S = A \bigcup B$ is not Radon's - a contradiction. $\square$

Corollary 4.1. *Let $q_1, \ldots, q_n$ be points on a convex curve K in $\mathbb{R}^{d-1}$. Let $p_i = (q_i, 1) \in \mathbb{R}^d$. Denote by $P(n, d)$ a convex polytope with vertices $\{p_1, -p_1, \ldots, p_n, -p_n\}$. Then the origin $0 \in P(n, d)$is covered by edges $(p_i, -p_i)$ and by d-simplices with vertices $p_{k_0}, -p_{k_1}, \ldots, (-1)^d p_{k_d}$ and $-p_{k_0}, p_{k_1}, \ldots, (-1)^{d+1} p_{k_d}$, where $1 \leq k_0 < k_1 < \ldots < k_d \leq n$.*

Proof. Clearly, $0 \in (-p_i, p_i)$. Let Δ be a simplex formed by vertices of $P(n, d)$. Let $V(\Delta) = A \bigcup (-B)$, where A and B are vertices with $x_d = 1$. It is easy to see that $0 \in \Delta$ if and only if $\operatorname{conv}(A) \bigcap \operatorname{conv}(B) = \emptyset$, i.e. $S = A \bigcup B$ is the Radon partition of S. Then Theorem 4.1 implies that A and B alternate along K. It proves the corollary. $\square$

If we apply Theorem 3.2 for $P = P(n, d)$, then Corollary 4.1 implies the following theorem.

Theorem 4.2. *Let M^d be a closed PL BUT manifold with a free involution. Let Λ be any equivariant triangulation of M. Let $L : V(\Lambda) \to \{+1, -1, +2, -2, \dots, +n, -n\}$ be an equivariant labelling. Then there is a complementary edge or an odd number of d-simplices whose labels are of the form $\{k_0, -k_1, k_2, \dots, (-1)^d k_d\}$, where $1 \le k_0 < k_1 < \dots < k_d \le n$.*

For the case $M = \mathbb{S}^d$ this is Fan's combinatorial lemma [4].

5 Sperner and Tucker's type lemmas for the case $d < m$

Now we consider extensions of the polytopal Sperner and Tucker lemmas for the case when $d = \dim P < \dim M = m$. In this case the set of fully-colored d-simplices defines certain $(m - d)$-submanifold S of M. Actually, a natural extension of Theorem 2.1 is that S is cobordant to zero. We also consider the Tucker lemma in terms of $\mathbb{Z}_2$-cobordisms.

An m-manifold M^m is called *null-cobordant* (or *cobordant to zero*) if there is a cobordism between M and the empty manifold; in other words, if M is the entire boundary of some $(m + 1)$-manifold. Equivalently, its cobordism class is trivial.

Definition. Let T be a triangulation of a closed PL manifold M^m. Let L be an n-labelling of T. Let C be a collection of n-labelled simplices of dimension $d \le m$. Now we define a simplicial complex $S(L, C)$.

Let V be the set of d-simplices of T which have the same labelling as simplices of C. We take V as the set of vertices of $S(L, C)$, i.e. $V(S(L, C)) := V$. All other simplices of $S(L, C)$ are defined by the following rule: a set of vertices $s_1, \dots, s_{k+1} \in V$ forms a k-simplex in $S(L, C)$ if simplices $s_1, \dots, s_{k+1} \in T$ have a common $(d - k)$-simplex $\sigma \in T$.

Lemma 5.1. *Let T be a triangulation of a closed PL-manifold M^m. Let L be an n-labelling of T. Then for any collection C of n-labelled simplices of dimension $d \le m$ the complex $S(L, C)$ is PL-homeomorphic to a subcomplex of M.*

In particular, if $P = \{p_1, \dots, p_n\} \subset \mathbb{R}^d$, $y \in \mathbb{R}^d$, $C = \operatorname{cov}_P(y)$, and $\operatorname{cov}_P(y)$ consists of d-simplices, then $S(L, C)$ is PL homeomorphic to $f_{L,P}^{-1}(y)$. Moreover, $S(L, \operatorname{cov}_P(y))$ is a closed manifold of dimension $m - d$.

Proof. The first part of the lemma follows from the following construction:
For any d-simplex $s_i \in V$ take an internal point v_i. Let the set of points $\{v_i\}$ be the vertices of a complex $Q \subset M$. If s_i and s_j from V have a common $(d - 1)$-simplex, then we join v_i and v_j by the edge e_{ij}. If three simplices s_i, s_j and s_k have a common $(d - 1)$-simplex, then

we add to Q a triangle (2-simplex) t_{ijk}. We can continue this process up to dimension $n-d$. It is clear that Q is PL-homeomorphic to $S(L,C)$.

Since $\mathrm{cov}_P(y)$ consists of d-simplices, the map $f_{L,P}: M \to \mathbb{R}^d$ is transversal to y. Then $R := f_{L,P}^{-1}(y)$ is a closed $(d-m)$-submanifold of M. If we take as the vertex v_i an internal point from $s_i \cap R$, then we obtain that Q and R are PL-homeomorphic. □

Theorem 5.1. *Let $P = \{p_1, \dots, p_n\}$ be a set in $\mathbb{R}^d$. Suppose $y \in \mathbb{R}^d$ is such that $\mathrm{cov}_P(y)$ consists of d-simplices. Then for any n-labelling L of a triangulation T of a closed PL-manifold M the manifold $S(L, \mathrm{cov}_P(y))$ is null-cobordant.*

Note that in the case $d = m$ the theorem yields that $S(L,C)$ consists of even number of points (see Theorem 2.1).

Proof. Let

$$W := M \times [0,1],\ M_0 := M \times \{0\},\ \text{and } M_1 := M \times \{1\}.$$

Let $f_0 := f_{L,P}: M_0 \to \mathbb{R}^d$ and $f_1(x) = q$ for any $x \in M_1$ where $q \neq y$ be a point in $\mathbb{R}^d$.

Note that f_0 and f_1 are PL maps transversal to y (or generic with respect to y). In other words, y is a regular image for f_i. It is well known that maps $f_i : M_i \to \mathbb{R}^d$ can be extended to a regular (transversal to y) PL map $F: W \to \mathbb{R}^d$ such that $F|_{M_0} = f_0$ and $F|_{M_1} = f_1$. (For instance, see our equivariant extension of this fact [11, Lemma 3].) Then $Z_F := F^{-1}(y)$ is a manifold of dimension $(n+1-d)$.

Denote $Z_i := Z_F \bigcap M_i = f_i^{-1}(y)$, $i = 0,1$. From Lemma 5.1 follows that Z_0 is PL-homeomorphic to $S(L, \mathrm{cov}_P(y))$. It is clear, Z_1 is empty. Thus, Z_0 is the boundary of Z_F, i.e. is null-cobordant. □

The case when P is the set of vertices of a d-simplex in $\mathbb{R}^d$ immediately implies

Corollary 5.1. *Let T be a triangulation of a closed PL-manifold M^m. Let L be an $(d+1)$-labelling of T, where $d \leq m$. Let C be a fully labelled d- simplex. Then $S(L,C)$ is a cobordant to zero.*

Let M be a closed PL-manifold with a free simplicial involution $T: M \to M$. We consider here antipodal (or equivariant) maps $f: M^m \to \mathbb{R}^d$, i.e. $f(T(x)) = -f(x)$, with $m < d$.

Now we extend the BUT manifolds.

Definition. We say that a closed PL free $\mathbb{Z}_2$-manifold (M^m, T) is a $\mathrm{BUT}_{m,d}$ if for any continuous $g: M^m \to \mathbb{R}^d$ there is a point $x \in M$ such that $g(T(x)) = g(x)$. Equivalently, if a continuous map $f: M^m \to \mathbb{R}^d$ is antipodal, then the zeros set $Z_f := f^{-1}(0)$ is not empty.

In our paper [11] we found a sufficient condition for (M^m, T) to be a $\mathrm{BUT}_{m,d}$. Let $\mathfrak{N}_*(\mathbb{Z}_2)$ denote the unoriented cobordism group of free involutions. Then $\mathfrak{N}_*(\mathbb{Z}_2)$ is a free $\mathfrak{N}_*$-module with basis $[\mathbb{S}^n, A]$, $n \geq 0$, where $[\mathbb{S}^n, A]$ is the cobordism class of the antipodal involution on the n-sphere. Thus, each $\mathbb{Z}_2$-manifold can be uniquely represented in $\mathfrak{N}_m(\mathbb{Z}_2)$ in the form:

$$[M,T] = \sum_{k=0}^{m} [V^k][\mathbb{S}^{m-k}, A].$$

We have the following theorem

Theorem 5.2 ([11],Corollary 5.2)**.** *Let* $[M^m, T] = [V^0][\mathbb{S}^m, A] + [V^1][\mathbb{S}^{m-1}, A] + \ldots + [V^m][\mathbb{S}^0, A]$ *in* $\mathfrak{N}_m(\mathbb{Z}_2)$ *with* $[V^{m-n}] \neq 0$ *in* $\mathfrak{N}_{m-n}$*, where* $m \geq n \geq d$*. Then for any antipodal continuous* $f : M^m \to \mathbb{R}^d$ *the set* Z_f *is not empty, i.e.* (M^m, T) *is* $BUT_{m,d}$

Any equivariant labelling L of a triangulation $\Lambda : V(\Lambda) \to \{+1, -1, +2, -2, \ldots, +d, -d\}$ of M defines a simplicial map $f_L : \Lambda \to C^d$, where C^d is the crosspolytope in $\mathbb{R}^d$ (see Section 3). It is easy to see that if L has no complementary edges, then $f_L : \Lambda \to \mathbb{R}^d$ has no zeros. It implies the following theorem.

Theorem 5.3. *Let* $m > d$*. Let* Λ *be any equivariant triangulation of a* $BUT_{m,d}$ *manifold* (M^m, T)*. Let*

$$L : V(\Lambda) \to \{+1, -1, +2, -2, \ldots, +d, -d\}$$

be any equivariant labelling of Λ*. Then there exists a complementary edge in* Λ*.*

This theorem is an extension of Theorem 3.1. It is not hard for the case $m > d$ to extend other theorems and corollaries from Sections 3 and 4.

Acknowledgment. I wish to thank Arseniy Akopyan and Frédéric Meunier for helpful discussions and comments.

References

[1] K. T. Atanassov, On Sperner's lemma, *Studia Sci. Math. Hungar.*, **32** (1996), 71-74.

[2] M. Breen, Primitive Radon partitions for cyclic polytopes, *Isr. J. Math.,* **15** (1973), 156-157.

[3] J. A. De Loera, E. Peterson, and F. E. Su, A Polytopal Generalization of Sperner's Lemma, *J. of Combin. Theory Ser. A,* **100** (2002), 1-26.

[4] K. Fan, A generalization of Tucker's combinatorial lemma with topological applications. *Ann. of Math.,* **56** (1952), 431-437.

[5] K. Fan, Simplicial maps from an orientable n-pseudomanifold into $\mathbf{S}^m$ with the octahedral triangulation, *J. of Combinatorial Theory,* **2** (1967), 588-602.

[6] J. Matoušek, Using the Borsuk-Ulam theorem, Springer-Verlag, Berlin, 2003.

[7] F. Meunier, Sperner labellings: a combinatorial approach, *J. of Combin. Theory Ser. A,* **113** (2006), 1462-1475.

[8] M. D. Meyerson and A. H. Wright, A new and constructive proof of the Borsuk-Ulam theorem, *Proc. Amer. Math. Soc.*, **73** (1979), 134-136.

[9] J. W. Milnor, Topology from the differentiable viewpoint, The University Press of Virginia, Charlottesville, Virginia, 1969.

[10] O. R. Musin, Chebyshev systems and zeros of a function on a convex curve, *Proc. Steklov Inst. of Math.,* **221** (1998), 236-246, arXiv:0903.1908.

[11] O. R. Musin, Borsuk-Ulam type theorems for manifolds, *Proc. Amer. Math. Soc.* **140** (2012), 2551-2560.

[12] T. Prescott and F. E. Su. A constructive proof of Ky Fan's generalization of Tucker's lemma, J. Combin. Theory Ser. A, **111** (2005), 257-265.

[13] E. Sperner, Neuer Beweis für die Invarianz der Dimensionszahl und des Gebietes, Abh. Math. Sem. Univ. Hamburg **6** (1928), 265-272.

[14] A. W. Tucker, Some topological properties of the disk and sphere. In: Proc. of the First Canadian Math. Congress, Montreal, 285-309, 1945.

[15] G. M. Ziegler, Lectures on Polytopes, Springer-Verlag, New York, 1995.

O. R. Musin, Department of Mathematics, University of Texas at Brownsville, 80 Fort Brown, Brownsville, TX, 78520.
E-mail address: oleg.musin@utb.edu